

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 875 512 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

04.11.1998 Bulletin 1998/45

(51) Int Cl.⁶: **C07D 403/06; A61K 31/495**(21) Application number: **98500085.0**(22) Date of filing: **07.04.1998**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

Designated Extension States:

AL LT LV MK RO SI

(72) Inventors:


- **Orjales Venero, Aurelio**
48990 Néguri (Vizcaya) (ES)
- **García Domínguez, Nefali**
48990 Getxo (Vizcaya) (ES)

(30) Priority: **16.04.1997 ES 9700812**(74) Representative: **Isern-Cuyas, María Luisa****Paseo de la Castellana****131, Bajo C****28046 Madrid (ES)**

(71) Applicant: **FABRICA ESPAÑOLA DE
PRODUCTOS
QUÍMICOS Y FARMACEUTICOS, S.A. (FAES)**
E-48940 Lejona-Lamiaco (Vizcaya) (ES)

(54) **New naphthylpiperazine derivatives with antipsychotic activity**

(57) A description is given of new naphthylpiperazine derivatives of general formula (I)



in which R_1 and R_2 , equal to or different from each other, can be hydrogen, a short chain alkyl group, halogen, a nitro group, an amino group, an acylamino group and a short chain alkoxy group; n can take values between 2 and 5; and R_3 can be a methoxy radical, a fluorine and a hydrogen atom, and their addition salts with pharmaceutically acceptable acids.

The compounds present affinity for serotonin $5HT_{1A}$ and $5HT_2$ receptors and for dopamine D_2 receptors, and can be useful as antipsychotics.

Their preparation is also described.

EP 0 875 512 A2

Description

Object of the Invention

The present invention relates to a number of new naphthylpiperazine derivatives and their addition salts, as well as their preparation process. The new compounds present serotonergic and central dopaminergic activity and are useful as antipsychotics.

Background of the Invention


The most widely accepted theory explaining the biochemical bases of schizophrenia holds that the dopaminergic activity in the mesolimb system of the brain is increased and, accordingly, the pharmacological power of classic antipsychotics is correlated with their affinity for D_2 receptors (Science 1976; 1982, 481-483). It has further been proposed (J. Pharm. Exp. Ther., 1989, 251, 238-246) that a high affinity for $5HT_2$ receptors in atypical antipsychotics would be responsible for the beneficial effects of the pharmacological profile of such atypical antipsychotics. It has moreover been found (J. Neurol. Transm., 1983, 57, 255) that $5HT_{1A}$ antagonists are capable of inverting haloperidol-induced catalepsy. Thus, compounds with affinity for $5HT_{1A}$, $5HT_2$ and D_2 receptors behave as atypical antipsychotics (Advances in Med. Chem., 3, 1995, 1-55).

European Patent EP 0329168 describes a number of 1,4-disubstituted piperazines with psychotropic activity in which the 1-substituent is a bicyclic heterocycle which incorporates an amide or imide function and the 4-substituent is a heterocycle.

US 4,983,606 discloses a number of compounds with anti-platelet aggregating, anti-vasospastic and anti-proliferative activities which carry the phthalazinone group bonded to a phenyl(benzoyl)piperazine or phenyl(benzoyl)piperidine fragment through a carbon chain.

J. Med. Chem. 1994, 37, 1060-1062 describes a number of new arylpiperazines having a high affinity for D_2 , D_3 , $5HT_{1A}$ and α_{1A} receptors, providing them with interesting antipsychotic properties, moreover presenting a low potential for causing extrapyramidal effects. Finally, Patent WO 93/16073 describes a number of arylpiperidines and arylpiperazines with D_2 and $5HT_2$ receptor antagonist properties, useful in the treatment of psychosis.

The present invention relates to a number of new naphthylpiperazine derivatives of formula (I) and their addition salts with pharmaceutically acceptable acids,




in which R_1 and R_2 (equal to or different from each other) can be hydrogen, a short chain alkyl group, halogen, a nitro group, an amino group, an acylamino group and a short chain alkoxy group; n can take values between 2 and 5, and R_3 can be a methoxyl radical, a hydrogen atom, and a fluorine atom.

The compounds described herein are essentially different from those referred to in the above-mentioned publications, due to the presence of the naphthylpiperazine fragment, which pharmacologically characterises them and provides them with a great affinity for $5HT_{1A}$ and $5HT_2$ receptors. These new compounds also have a great affinity for the D_2 receptor and are therefore useful as atypical antipsychotics.


Description of the Invention

The preparation of compounds of formula (I) can be carried out along various synthetic routes, using conventional methods:

a) Condensation of a compound of formula (II)



in which R_1 and R_2 have the meaning specified for formula (I) and Z is chlorine or bromine, with a piperazine of formula (III)




The reaction is carried out in an inert solvent, in the presence of a base, and at a temperature ranging between 20°C and the boiling point of the reaction mixture.

Examples of used solvents are dichloromethane, chloroform, acetonitrile, dimethylformamide and tetrahydrofuran.

Used bases can be an alkaline carbonate or bicarbonate, such as K_2CO_3 , $KHCO_3$, Na_2CO_3 , Na_2HCO_3 or a tertiary amine such as pyridine or triethylamine.


The reaction rate may be increased by adding catalytic amounts of an alkaline iodide, such as KI to the reaction mixture.

Piperazines of formula (III) are prepared by reacting bis (2-chloroethyl)amine with the corresponding naphthylamine. Compounds of general formula (II) used as starting products at the previous condensation stage are prepared by reacting a compound of general formula (IV)



in which R_1 and R_2 have been previously defined, with an alkyl dihalide in the presence of a strong base, such as NaH in an aprotic solvent such as dimethylformamide or tetrahydrofuran, or solid potassium hydroxide in dimethylformamide, at a temperature ranging between room temperature and 100°C. Compounds of general formula (IV) are prepared by reacting suitable 2-formylbenzoic acids with hydrazine hydrate.

Alternately, compounds of general formula (II) can be prepared by substitution of the hydroxyl group in compounds of formula (V) by chlorine or bromine.



The substitution reaction can be carried out by treatment with hydrogen chloride, hydrogen bromide, or by treatment

with organic or inorganic acid halides, such as thionyl chloride, oxalyl chloride, in an inert solvent such as chloroform, dichloromethane, or toluene, or using the acid chloride as a solvent, at a temperature ranging between 0°C and the boiling point of the reaction mixture.

Compounds of general formula (V) can in turn be prepared as specified in *Chimie Thérapeutique*, 1967, 2, 250-253.

As noted hereinbefore, the new compounds of general formula (I) show SNC activity, specifically at the 5HT_{1A}, 5HT₂ and D₂ receptor levels.

PHARMACOLOGICAL RECEPTOR BINDING STUDIES.

Binding studies have been carried out to ascertain the affinity of the compounds for 5HT_{1A}, 5HT₂ and D₂ receptors.

5HT_{1A} RECEPTOR

Cerebral cortex taken from both male and female Wistar rats was homogenised in saccharose buffer 0.32 M (1:10 g/mL) and centrifuged at 900 g (10 min, 4°C). The supernatant was collected and centrifuged at 48000 g (25 min, 4°C). The sediment thus obtained was resuspended in cold TRIS buffer 50 mM (pH 7.5, 1:10, g/mL), homogenised, incubated at 37°C for 15 min and centrifuged again at 48000 g (25 min, 4°C). The final sediment was re-suspended in cold SNAYDER buffer (1:4, g/mL), homogenised and kept at -70°C in 5 mL containers.

For the displacement trial, 100 µL of radioligand (2 nM, final conc.), 100 µL of the different tested concentrations of the displacing product and 750 µL of a suspension of membranes 1:32 in SNAYDER with pargilin were used. The volume was topped up to 1 mL with 50 µL of SNAYDER. Serotonin (5HT) 10 µM was used to define the non-specific binding.

5HT₂ RECEPTOR

Prefrontal cortex taken from both male and female Wistar rats was homogenised in saccharose buffer 0.25 M (1:10 g/mL) and centrifuged at 1080 g (10 min, 4°C). The supernatant was set aside and the pellet re-suspended in the same buffer (1:5, g/mL) and centrifuged again under the same conditions. The mixture of both supernatants was topped up to 1:40 (g/mL) with TRIS buffer 50 mM (pH 7.7) and centrifuged at 35000 g (10 min, 4°C). The sediment thus obtained was re-suspended in cold TRIS buffer (1:40, g/mL) and centrifuged again at 35000 g (10 min, 4°C). The final sediment was re-suspended in cold TRIS buffer (1:10, g/mL), homogenised and kept at -70°C in 5 mL containers.

For the displacement trial, 100 µL of radioligand (0.5-1 nM, final concentration), 100 µL of the different tested concentrations of the displacing product and 750 µL of a suspension of membranes 1:50 (0.54 mg/mL, 15 mg fresh tissue) in TRIS were used. The volume was topped up to 1 mL with 50 µL of TRIS/10% ethanol. Metisergide 1 µM was used to define the non-specific binding. The displacer, radioligand and Metisergide dilutions were all made with TRIS buffer with 10% ethanol (v/v).

D₂ RECEPTOR.

Corpus striatum taken from both male and female Wistar rats was homogenised in TRIS buffer 50 mM (pH 7.7, 1:50 g/mL) and centrifuged at 47800 g (10 min, 4°C). The supernatant was eliminated and the pellet re-suspended in the same buffer (1:50, g/mL), incubated at 37°C for 10 min and centrifuged again under the same conditions. The final sediment thus obtained was re-suspended in cold TRIS buffer 50 mM (pH 7.4, 1:10, g/mL) containing NaCl 120 mM + KCl 5 mM + CaCl₂ 2 mM + MgCl₂ 1 mM + ascorbic acid 0.01% g/mL, and kept at -70°C in 2.5 mL aliquots. Membrane dilutions (1:100-1:300) were subsequently carried out and the amount of proteins was assessed by the Lowry method.

For the displacement test, 100 µL of radioligand (1-2 nM, final conc.), 100 µL of the different tested concentrations of the displacing product and 750 µL of a suspension of membranes 1:150 (0.39-0.13 protein mg/mL) in the previous (saline) TRIS + 10 µM of pargilin were used. The volume was topped up to 1 mL with 50 µL of the previous TRIS. Butachlamol 1 µM was used to define the non-specific binding, which was added (100 µL) to the blank series. The displacer, radioligand and Butachlamol dilutions were all made with TRIS (saline) buffer + pargilin. The samples were incubated for 60 min at 25°C.

The products described in the present invention have shown a high (nanomolar range) affinity for the three tested receptors, which renders them potentially useful as antipsychotics.

The following examples provide further details of the invention, which is not howsoever limited to such examples.

Example 1.2-(4-Bromobutyl)-1(2H)-phthalazinone.

Small portions (2.5 g, 60 mmol) of a dispersion of 60% NaH in mineral oil were added over a solution of 1(2H)-phthalazinone (7.3 g, 50 mmol) in dimethylformamide (100 mL) cooled to 0°C. The reaction mixture was left to reach room temperature and kept stirred for an hour. 1,4-dibromobutane (18 mL, 150 mmol) was then added at a time and the reaction mixture was stirred, at room temperature, for 4 hours. The reaction mixture was poured over crushed ice, extracted with ethyl ether (twice), and the organic extracts were dried over anhydrous Na₂SO₄ and concentrated to dryness. The excess dibromobutane was eliminated by distillation and the obtained residue was purified by flash chromatography (CH₂Cl₂), yielding an oil (11.6 g, yield: 83%) identified on the basis of its spectroscopic data as the title product.

Example 2.2-[4-(4-naphthylpiperazine-1-yl)-butyl]-1(2H)-phthalazinone.

A mixture of 2-(4-bromobutyl)-1(2H)-phthalazinone (1.7 g, 6 mmol), 1-naphthylpiperazine (1.3 g, 6 mmol), K₂CO₃ (0.84 g, 6 mmol), and KI (10 mg) in acetonitrile (50 mL) was refluxed for 4 hours. When heating was over, the solvent was eliminated at reduced pressure and the residue distributed between dichloromethane and water; the aqueous phase was extracted with dichloromethane (twice). The organic extracts were collected, dried over anhydrous Na₂SO₄ and concentrated to dryness. The obtained residue was purified by flash column (CH₂Cl₂/MeOH 97:3), yielding 2-[4-(4-naphthylpiperazine-1-yl)-butyl]-1(2H)-phthalazinone (1.7 g). Treatment with hydrochloric ethanol yielded a white solid, which was re-crystallised in methanol, yielding the monohydrochloride of the title compound, presenting a M.P. higher than 260°C.

Example 3.2-(2-hydroxyethyl)-6,7-dimethoxy-1(2H)-phthalazinone.

2-hydroxyethylhydrazine (4.2 g, 55 mmol) was added to a solution of 6-formyl-3,4-dimethoxybenzoic acid (10.5 g, 50 mmol) in ethanol (100 mL). The reaction mixture was refluxed for 3 hours, after which the solvent was eliminated at reduced pressure. The resultant residue was treated with ethyl ether to yield a white solid which was collected by filtration, yielding 2-(2-hydroxyethyl)-6,7-dimethoxy-1(2H)-phthalazinone (9.2 g, yield: 76%). M.P. = 176-179°C.

Example 4.2-(2-chloroethyl)-6,7-dimethoxy-1(2H)-phthalazinone.


Thionyl chloride (15 mmol) was added to a solution of 2-[2-(hydroxyethyl)]-6,7-dimethoxy-1(2H)-phthalazinone (2.5 g, 10 mmol) in chloroform (50 mL) and the mixture was kept stirred at room temperature overnight. The solvent was eliminated at reduced pressure, and the obtained solid was treated with hexane, filtered and purified by flash chromatography (CH₂Cl₂), yielding 2-(2-chloroethyl)-6,7-dimethoxy-1(2H)-phthalazinone (2.2 g, yield: 81%). M.P. 178-181°C.

Example 5.2-[2-(4-naphthylpiperazine-1-yl)-ethyl]-6,7-dimethoxy-1(2H)-phthalazinone.

A mixture of 2-(2-chloroethyl)-6,7-dimethoxy-1(2H)-phthalazinone (1.6 g, 6 mmol), 1-naphthylpiperazine (1.3 g, 6 mmol), K₂CO₃ (0.84 g, 6 mmol) and KI (10 mg) in acetonitrile (50 mL) was refluxed for 7 hours. When heating was over, the solvent was eliminated at reduced pressure and distributed between dichloromethane and water, and the aqueous phase was extracted with dichloromethane (twice). The organic extracts were collected, dried over anhydrous Na₂SO₄ and concentrated to dryness. The obtained residue was purified by flash chromatography (CH₂Cl₂/MeOH 97:3) yielding 2-[2-(4-naphthylpiperazine-1-yl)-ethyl]-6,7-dimethoxy-1(2H)-phthalazinone (1.5 g). Treatment with hydrochloric ethanol yielded a white solid, which was re-crystallised in methanol, yielding the monohydrochlorate of the title compound, presenting a M.P. higher than 260°C.

Claims

1. New naphthylpiperazine derivatives of general formula (I)



in which R_1 and R_2 , equal to or different from each other, are hydrogen, a short chain alkyl group, halogen, a nitro group, an amino group, an acylamino group and a short chain alkoxy group; n can take values between 2 and 5; and R_3 can be a methoxyl radical, a fluorine and a hydrogen atom.

2. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[4-(4-naphthylpiperazine-1-yl)-butyl]-1(2H)-phthalazinone.
3. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[3-(4-naphthylpiperazine-1-yl)-propyl]-1(2H)-phthalazinone.
4. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[2-(4-naphthylpiperazine-1-yl)-ethyl]-1(2H)-phthalazinone.
5. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[4-(4-naphthylpiperazine-1-yl)-butyl]-6,7-dimethoxy-1(2H)-phthalazinone.
6. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[2-(4-naphthylpiperazine-1-yl)-ethyl]-6,7-dimethoxy-1(2H)-phthalazinone.
7. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[4-(4-naphthylpiperazine-1-yl)-butyl]-7-nitro-1(2H)-phthalazinone.
8. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[4-(4-naphthylpiperazine-1-yl)-butyl]-7-amino-1(2H)-phthalazinone.
9. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[4-(4-naphthylpiperazine-1-yl)-butyl]-7-acetylamino-1(2H)-phthalazinone.
10. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[3-(4-naphthylpiperazine-1-yl)-propyl]-7-nitro-1(2H)-phthalazinone.
11. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[4-(4-(7-methoxynaphthyl)-piperazine-1-yl)-butyl]-1(2H)-phthalazinone.
12. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[3-(4-(7-methoxynaphthyl)-piperazine-1-yl)-propyl]-1(2H)-phthalazinone.
13. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[4-(4-(7-methoxynaphthyl)-piperazine-1-yl)-butyl]-7-nitro-1(2H)-phthalazinone.
14. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[3-(4-(7-methoxynaphthyl)-piperazine-1-yl)-propyl]-7-nitro-1(2H)-phthalazinone.
15. New naphthylpiperazine derivatives, in accordance with claim 1, characterised by comprising a compound consisting of 2-[4-(4-(7-methoxynaphthyl)-piperazine-1-yl)-butyl]-6,7-dimethoxy-1(2H)-phthalazinone.

16. New naphthylpiperazine derivatives, in accordance with claims 1 to 15, characterised by comprising addition salts with pharmaceutically acceptable acids of the compounds of formula (I).

17. New naphthylpiperazine derivatives, in accordance with claims 1 to 15, characterised by the use of the compounds of formula (1) as antipsychotics.

5

10

15

20

25

30

35

40

45

50

55

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 875 512 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
07.04.1999 Bulletin 1999/14

(51) Int Cl.⁶: C07D 403/06, A61K 31/495

(43) Date of publication A2:
04.11.1998 Bulletin 1998/45

(21) Application number: 98500085.0

(22) Date of filing: 07.04.1998

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• Orjales Venero, Aurelio
48990 Neguri (Vizcaya) (ES)
• Garcia Dominguez, Neftali
48990 Getxo (Vizcaya) (ES)


(30) Priority: 16.04.1997 ES 9700812

(74) Representative: Isern-Cuyas, Maria Luisa
Paseo de la Castellana
131, Bajo C.
28046 Madrid (ES)

(71) Applicant: FABRICA ESPANOLA DE
PRODUCTOS
QUIMICOS Y FARMACEUTICOS, S.A. (FAES)
E-48940 Lejona-Lamiaco (Vizcaya) (ES)

(54) New naphthylpiperazine derivatives with antipsychotic activity

(57) A description is given of new naphthylpiperazine derivatives of general formula (I)



in which R₁ and R₂, equal to or different from each other, can be hydrogen, a short chain alkyl group, halogen, a nitro group, an amino group, an acylamino group and a short chain alkoxy group; n can take values between 2 and 5; and R₃ can be a methoxyl radical, a fluorine and a hydrogen atom, and their addition salts with pharmaceutically acceptable acids.

The compounds present affinity for serotonin 5HT_{1A} and 5HT₂ receptors and for dopamine D₂ receptors, and can be useful as antipsychotics.

Their preparation is also described.

EP 0 875 512 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 50 0085

DOCUMENTS CONSIDERED TO BE RELEVANT					
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)		
Y	WO 96 16949 A (PF MEDICAMENT ; PATOISEAU JEAN FRANCOIS (FR); FAURE CHRISTIAN (FR);) 6 June 1996 * see example 10, page 9 * * the whole document *	1-17	C07D403/06 A61K31/495		
Y,D	EP 0 329 168 A (BRISTOL MYERS CO) 23 August 1989 * the whole document *	1-17			
Y	WO 95 01965 A (PF MEDICAMENT ; PATOISEAU JEAN FRANCOIS (FR); COURET FRANCOISE (FR)) 19 January 1995 * the whole document *	1-17			
Y	EP 0 527 081 A (PF MEDICAMENT) 10 February 1993 * the whole document *	1-17			
Y	DUKIC, S. ET AL.: "Structure-activity Relationship Studies on D-2/5-HT1A Receptor Ligands 4-(2-Heteroarylethyl)-1-arylpiperazines " ARZNEIM.FORSCH./DRUG RES., vol. 47, no. 3, 1997, pages 239-243, XP002092306 HEIDELBERG * the whole document *	1-17			
Y	MELTZER, H.Y. ET AL.: "Classification of Typical and Atypical Antipsychotic Drugs on the Basis of Dopamine D-1, D-2, and Serotonin2 pki Values " J. PHARMACOL. EXP. THER., vol. 25, no. 1, 1989, pages 238-246, XP002092307 BALTIMORE * the whole document *	1-17	<table border="1"> <thead> <tr> <th>TECHNICAL FIELDS SEARCHED (Int.Cl.6)</th> </tr> </thead> <tbody> <tr> <td>C07D</td> </tr> </tbody> </table>	TECHNICAL FIELDS SEARCHED (Int.Cl.6)	C07D
TECHNICAL FIELDS SEARCHED (Int.Cl.6)					
C07D					
The present search report has been drawn up for all claims					
Place of search MUNICH		Date of completion of the search 5 February 1999	Examiner Stellmach, J		
<table border="0"> <tr> <td> CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document </td> <td> T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons & : member of the same patent family, corresponding document </td> </tr> </table>				CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document	T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons & : member of the same patent family, corresponding document
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document	T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons & : member of the same patent family, corresponding document				

EPO FORM 1503 03/92 (P04001)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 50 0085

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	EP 0 434 561 A (ADIR) 26 June 1991 * the whole document *	1-17	
A,D	US 4 983 606 A (CASAGRANDE CESARE ET AL) 8 January 1991 * the whole document *	1-17	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 5 February 1999	Examiner Stellmach, J
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.92 (PUBO1)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 50 0085

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-02-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9616949 A	06-06-1996	FR 2727682 A	07-06-1996
		AT 169008 T	15-08-1998
		AU 4306796 A	19-06-1996
		CA 2206794 A	06-06-1996
		DE 69503803 D	03-09-1998
		EP 0794949 A	17-09-1997
		ES 2121441 T	16-11-1998
		JP 10510286 T	06-10-1998
		NZ 297822 A	24-09-1998
EP 0329168 A	23-08-1989	US 5001130 A	19-03-1991
		JP 1249769 A	05-10-1989
		US 5116970 A	26-05-1992
		US 5206365 A	27-04-1993
WO 9501965 A	19-01-1995	FR 2707294 A	13-01-1995
		AT 152445 T	15-05-1997
		AU 679761 B	10-07-1997
		AU 7127794 A	06-02-1995
		CA 2166629 A	19-01-1995
		DE 69402981 D	05-06-1997
		EP 0707571 A	24-04-1996
		JP 8512304 T	24-12-1996
		NZ 268384 A	27-08-1996
		US 5591743 A	07-01-1997
EP 0527081 A	10-02-1993	FR 2680172 A	12-02-1993
		AT 132494 T	15-01-1996
		DE 69207301 D	15-02-1996
		DE 69207301 T	29-08-1996
		DK 527081 T	09-04-1996
		ES 2083708 T	16-04-1996
		GR 3019274 T	30-06-1996
		JP 5194441 A	03-08-1993
		US 5308844 A	03-05-1994
EP 0434561 A	26-06-1991	FR 2655988 A	21-06-1991
		AT 129241 T	15-11-1995
		AU 635369 B	18-03-1993
		AU 6823590 A	27-06-1991
		CA 2032713 A	21-06-1991
		DE 69023105 D	23-11-1995
		DE 69023105 T	30-05-1996
		DK 434561 T	04-03-1996
		ES 2080815 T	16-02-1996
		GR 3018472 T	31-03-1996

EPO FORM P0689

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 98 50 0085

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-02-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0434561 A		IE 78927 B	15-01-1997
		JP 1942670 C	23-06-1995
		JP 3291275 A	20-12-1991
		JP 6076395 B	28-09-1994
		OA 9477 A	15-11-1992
		PT 96255 A,B	30-09-1991
		US 5143916 A	01-09-1992
		US 5166157 A	24-11-1992
		US 5162324 A	10-11-1992
		US 5162321 A	10-11-1992
		US 5166156 A	24-11-1992
US 4983606 A	08-01-1991	JP 1151561 A	14-06-1989

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82